

MODULI SPACES OF J -HOLOMORPHIC CURVES WITH GENERAL JET CONSTRAINTS

KE ZHU

ABSTRACT. In this paper, we prove that the tangent map of the holomorphic k -jet evaluation j_{hol}^k from the mapping space to holomorphic k -jet bundle, when restricted on the universal moduli space $\mathcal{M}_1^*(\Sigma, M, \beta)$ of simple J -holomorphic curves with one marked point, is surjective. From this we derive that for generic J , the moduli space of simple J -holomorphic curves with general jet constraints at marked points is a smooth manifold of expected dimension.

1. INTRODUCTION

Let (M, ω) be a symplectic manifold of dimension $2n$. Denote by $\mathcal{J}_\omega$ the set of almost complex structures J on M compatible with ω . Let Σ be a compact oriented surface without boundary, and (j, u) a pair of complex structure j on Σ and a map $u : \Sigma \rightarrow M$. We say (j, u) is a J -holomorphic curve if $\bar{\partial}_{J,j}u := \frac{1}{2}(du + J \circ du \circ j) = 0$. We let $\mathcal{M}_1(\Sigma, M, J; \beta)$ be the standard moduli space of J -holomorphic curves in class $\beta \in H_2(M, \mathbb{Z})$ with one marked point, and $\mathcal{M}_1^*(\Sigma, M, J; \beta)$ be the set of simple (i.e. somewhere injective) J -holomorphic curves in $\mathcal{M}_1(\Sigma, M; \beta)$.

Since the birth of the theory of J -holomorphic curves, moduli spaces of J -holomorphic curves with constraints at marked points have lead to finer symplectic invariants like Gromov-Witten invariants and quantum cohomology. J -holomorphic curves with embedding property also plays important role in low dimensional symplectic geometry, like the works of [HT] and [Wen]. These constraints all can be viewed as partial differential relations in the 0-jet and 1-jet bundles. In relative Gromov-Witten theory, contact order of J -holomorphic curves with given symplectic hypersurfaces (divisors) was used to define the relevant moduli spaces. In the work of Cieliebak-Mohke [CM] and Oh [Oh], the authors studied the moduli space of J -holomorphic curves with prescribed vanishing orders of derivatives at marked points. All these are vanishing conditions in k -jets bundles. It is then natural to ask what properties we can expect for moduli spaces of J -holomorphic curves with general constraints in jet bundles (while all constraints in previous examples are *zero sections* in various jet bundles).

The main purpose of this paper is to confirm that for a wide class of closed partial differential relations in *holomorphic jet bundles* (Definition 1, originally defined in [Oh]), the moduli spaces of J -holomorphic curves from Σ to M with given constraints at marked points behave well for generic J (Theorem 3). Namely, they are smooth manifolds of dimension predicted by index theorem, and all elements in

Date: November 9, 2009.

Key words and phrases. J -holomorphic curve, jet evaluation map, transversality.

the moduli spaces are Fredholm regular. During the proof it appears that holomorphic jet bundles are the natural framework to put jet constraints for J -holomorphic curves in order to obtain regularity of their moduli spaces. The regularity of J -holomorphic curve moduli spaces *fails* for general constraints in *usual* jet bundle (Remark 2), but still holds in a special case when the moduli space consists of immersed J -holomorphic curves (Theorem 2).

The key of the proof is to establish the surjective property of the linearization of k -jet evaluations on the universal moduli spaces of J -holomorphic curves at marked points insider the mapping space, including the parameter $J \in \mathcal{J}_\omega$ (Theorem 1). It is important to take the evaluations in *holomorphic jet bundles* in order to get the surjectivity of the linearization of the k -jet evaluation map.

Since $\mathcal{J}_\omega$ is a huge parameter space to deform J -holomorphic curves, the surjective property here is a reminiscence of the classic Thom transversality theorem, which says that the k -jet evaluation on smooth mapping space to the k -jet bundle is transversal to any section there.

The framework of the paper is similar to [Oh], which in turn is a higher jet generalization of [OZ] for 1-jet transversality of J -holomorphic curves. The main steps of the paper are in order:

- (1) We set up the Banach bundle including the finite dimensional holomorphic k -jet subbundle $J_{hol}^k(\Sigma, M)$ over the mapping space $\mathcal{F}_1(\Sigma, M) \times \mathcal{J}_\omega$ and define the section $\Upsilon_k = (\bar{\partial}, j_{hol}^k)$, where

$$\Upsilon_k : ((u, j, z_0), J) \rightarrow (\bar{\partial}_{j,J} u, j_{hol}^k(u(z_0))).$$

We inteprete the universal J -holomorphic curve moduli space as

$$\mathcal{M}(\Sigma, M) = \bar{\partial}^{-1}(0) = \Upsilon_k^{-1}(0, J_{hol}^k(\Sigma, M)).$$

- (2) We compute the linearization $D\Upsilon_k$ of the section Υ_k . We express the submersion property of Υ_k as the solvability of a system of equations $D\Upsilon_k(\xi, B) = (\gamma, \alpha)$ for any (γ, α) , where $(\xi, B) \in T_u \mathcal{F}_1(\Sigma, M) \times T_J \mathcal{J}_\omega$, or equivalently, the vanishing of the cokernal element (η, ζ) in the Fredholm alternative system: $F \langle (\xi, B), (\eta, \zeta) \rangle = 0$ for all (ξ, B) . This is called the cokernal equation.
- (3) Using the abundance of $B \in T_J \mathcal{J}_\omega$ we get $\text{supp } \eta \subset \{z_0\}$. Then we use a structure theorem in distribution to write η as a linear combination of δ function and its derivatives at z_0 , up to $(k-1)$ -th order derivatives.
- (4) Since $\text{supp } \eta \subset \{z_0\}$ the cokernal equation is supported at z_0 . We replace the ξ in the cokernal equation by $\xi + h$ where $h = h(z, \bar{z})$ is a suitable polynomial in local coordinates nearby z_0 , and set $B = 0$, so that the cokernal equation is reduced to $\langle D_u \bar{\partial}_{j,J} \xi, \eta \rangle = 0$ for all ξ . The crucial observation is that to get $\langle D_u \bar{\partial}_{j,J} \xi, \eta \rangle = 0$ we do not need so strong conditions of vanishing of $1 \sim k$ -derivatives of u at z_0 as in [Oh] and [CM]. This is by exploring the flexibility of h to get rid of redundant terms from the original cokernal equation.
- (5) Then we apply elliptic regularity to conclude $\eta = 0$ and consequently $\zeta = 0$. Therefore we get the surjectivity of $D\Upsilon_k$ and Dj_{hol}^k .
- (6) Finally, there is an obstruction in step 4 to get h when $\zeta_k = 0$, where ζ_k is the k -th component of ζ . But when $\zeta_k = 0$ the cokernal equation is

reduced to the $(k-1)$ -jet evaluation setting, so we still get $(\eta, \zeta) = (0, 0)$ by induction on k .

Acknowledgement. The author would like to thank Yakov Eliashberg to suggest the generalization from [Oh] to general PDE relations. He would also like to thank Yong-Geun Oh on past discussions in holomorphic jet transversality.

2. HOLOMORPHIC JET BUNDLE

We recall the holomorphic jet bundle from [Oh]. Given Σ, M , and $(z, x) \in \Sigma \times M$, the k -jet with source z and target x is defined as (see [Hir])

$$J_{z,x}^k(\Sigma, M) = \prod_{l=0}^k \text{Sym}^l(T_z \Sigma, T_x M),$$

where $\text{Sym}^l(T_z \Sigma, T_x M)$ is the set of l -multilinear maps from $T_z \Sigma$ to $T_x M$ for $l \geq 1$. Here for convenience we have set $\text{Sym}^0(T_z \Sigma, T_x M) = M$. Let

$$J^k(\Sigma, M) = \bigcup_{(z,x) \in \Sigma \times M} J_{z,x}^k(\Sigma, M)$$

be the k -jet bundle over $\Sigma \times M$. For the mapping space

$$\mathcal{F}_1(\Sigma, M; \beta) = \{((\Sigma, j), u) \mid j \in \mathcal{M}(\Sigma), z \in \Sigma, u : \Sigma \rightarrow M, [u] = \beta\},$$

we consider the map

$$\mathcal{F}_1(\Sigma, M; \beta) \rightarrow \Sigma \times M, (u, j, z) \rightarrow (z, u(z)).$$

By this map we can pull back the bundle $J^k(\Sigma, M) \rightarrow \Sigma \times M$ to the base $\mathcal{F}_1(\Sigma, M; \beta)$. By abusing notation, we still call the resulted bundle by $J^k(\Sigma, M)$. Then $J^k(\Sigma, M) \rightarrow \mathcal{F}_1(\Sigma, M; \beta)$ is a finite dimensional vector bundle over the Banach manifold $\mathcal{F}_1(\Sigma, M; \beta)$. We define the k -jet evaluation

$$j^k : \mathcal{F}_1(\Sigma, M; \beta) \rightarrow J^k(\Sigma, M), j^k((u, j), z) = j_z^k u \in J_{z, u(z)}^k(\Sigma, M).$$

Then j^k is a smooth section. Classic Thom transversality theorem says that j^k is transversal to any section in $J^k(\Sigma, M)$.

Now we turn to the case when Σ and M are equipped with (almost) complex structures j and J respectively. The corresponding concept is the holomorphic jet bundle defined in [Oh]. With respect to (j_z, J_x) , $\text{Sym}_{z,x}^l(\Sigma, M)$ splits into summands indexed by the bigrading (p, q) for $p + q = l$:

$$\text{Sym}_{z,x}^l(\Sigma, M) = \text{Sym}^{(l,0)}(T_z \Sigma, T_x M) \oplus \text{Sym}^{(0,l)}(T_z \Sigma, T_x M) \oplus \text{“mixed parts”}$$

Let

$$\begin{aligned} H_{j_z, J_x}^{(l,0)}(\Sigma, M) &= \text{Sym}^{(l,0)}(T_z \Sigma, T_x M), \\ H_{j,J}^{(l,0)}(\Sigma, M) &= \bigcup_{(z,x) \in \Sigma \times M} H_{j_z, J_x}^{(l,0)}(\Sigma, M). \end{aligned}$$

Given (j, J) , the (j, J) -holomorphic jet bundle $J_{(j,J)\text{hol}}^k(\Sigma, M)$ is defined as

$$(2.1) \quad J_{(j,J)\text{hol}}^k(\Sigma, M) = \prod_{l=0}^k H_{j,J}^{(l,0)}(\Sigma, M),$$

which is a finite dimensional vector bundle over $\Sigma \times M$.

We define the bundle

$$J_{hol}^k(\Sigma, M) = \bigcup_{(j,J) \in \mathcal{M}(\Sigma) \times \mathcal{J}_\omega} J_{(j,J)hol}^k(\Sigma, M).$$

$J_{hol}^k(\Sigma, M) \rightarrow \Sigma \times M \times \mathcal{M}(\Sigma) \times \mathcal{J}_\omega$ is a finite dimensional vector bundle over the base Banach manifold. Using the pull back of the map

$ev : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow \Sigma \times M \times \mathcal{M}(\Sigma) \times \mathcal{J}_\omega, ((u, j), z, J) \rightarrow (z, u(z), j, J),$
 $ev^*(J_{hol}^k(\Sigma, M))$ is a finite dimensional vector bundle over the Banach manifold $\mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega$. By abusing of notation, we still call $ev^*(J_{hol}^k(\Sigma, M))$ by $J_{hol}^k(\Sigma, M)$.

Definition 1. $J_{hol}^k(\Sigma, M) \rightarrow \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega$ is called the holomorphic k -jet bundle.

Let $\pi_{hol} : J_{hol}^k(\Sigma, M) \rightarrow J_{hol}^k(\Sigma, M)$ be the bundle projection. We define the holomorphic k -jet evaluation

$$j_{hol}^k = \pi_{hol} \circ j^k.$$

It is not hard to see j_{hol}^k is a smooth section of the Banach bundle $\mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow J_{hol}^k(\Sigma, M)$. According to the summand (2.1), we write j_{hol}^k in components

$$j_{hol}^k = \prod_{l=0}^k \sigma^l,$$

where the l -th component is

$$\sigma^l : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow H_{j,J}^{(l,0)}(\Sigma, M), ((u, j), z, J) \rightarrow \pi_{j,J}^{hol}(d^l u(z)).$$

We remark that if J is integrable, σ^l corresponds to the l -th holomorphic derivative $\frac{\partial^l}{\partial z^l} u$ of u at z .

The important point is that the holomorphic k -jet bundle and the section j_{hol}^k are canonically associated to the pair (Σ, j) and (M, J) in the “off-shell level”, i.e. on the space of all smooth maps, not only J -holomorphic maps. This enables us to formulate the jet constraints for J -holomorphic maps as some submanifold in the bundle $J_{hol}^k(\Sigma, M) \rightarrow \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega$.

3. FREDHOLM SET UP

The Fredholm set up is the same as in [Oh], with the simplification that we only need one marked point on Σ . The case with more marked points has no essential difference. We introduce the standard bundle

$$\mathcal{H}'' = \bigcup_{((u,j),J)} \mathcal{H}_{((u,j),J)}'', \quad \mathcal{H}_{((u,j),J)}'' = \Omega_{j,J}^{(0,1)}(u^* TM)$$

and define the section

$$\Upsilon_k : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow \mathcal{H}'' \times J^k(\Sigma, M)$$

as

$$\Upsilon_k((u, j), z, J) = (\bar{\partial}(u, j, J); j_{hol}^k(u, j, J, z)),$$

where

$$\bar{\partial}(u, j, J) = \bar{\partial}_{j,J}(u) = \frac{du + J \circ du \circ j}{2}.$$

Given $\beta \in H_2(M, \mathbb{Z})$, let

$$\mathcal{M}_1(\Sigma, M; \beta) = \bigcup_{J \in \mathcal{J}_\omega} \mathcal{M}_1(\Sigma, M, J; \beta)$$

be the universal moduli space of J -holomorphic curves in class β with one marked point. Its open subset consisting of somewhere injective J -holomorphic curves is denoted by $\mathcal{M}_1^*(\Sigma, M; \beta)$. It is a standard fact in symplectic geometry that $\mathcal{M}_1^*(\Sigma, M; \beta)$ is a Banach manifold.

Now we make precise the necessary regularity requirement for the Banach manifold set-up:

- (1) To make sense of the evaluation of $j^k u$ at a point z on Σ , we need to take at least $W^{k+1,p}$ -completion with $p > 2$ of $\mathcal{F}_1(\Sigma, M; \beta)$ so $j^k u \in W^{1,p} \hookrightarrow C^0$. To make the section Υ_k differentiable we need to take $W^{k+2,p}$ completion, since in (4.2) $(k+1)$ -th derivatives of u are involved. To apply Sard-Smale theorem, we actually need to take $W^{N,p}$ completion with sufficiently large $N = N(\beta, k)$.
- (2) We provide $\mathcal{H}''$ with topology of a $W^{N,p}$ Banach bundle.
- (3) We also need to provide the Banach manifold structure of $\mathcal{J}_\omega$. We can borrow Floer's scheme [F, F] for this whose details we refer readers thereto.

4. TRANSVERSALITY

Theorem 1. *At every J -holomorphic curve $((u, j), z, J) \in \mathcal{M}_1^*(\Sigma, M; \beta) \subset \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega$, the linearization $D\Upsilon_k$ of the map*

$$\Upsilon_k = (\bar{\partial}, j_{hol}^k) : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow \mathcal{H}'' \times J_{hol}^k(\Sigma, M)$$

is surjective. Especially the linearization Dj_{hol}^k of the holomorphic k -jet evaluation

$$j_{hol}^k : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow J_{hol}^k(\Sigma, M)$$

on $\mathcal{M}_1^(\Sigma, M; \beta)$ is surjective.*

To prove theorem we need to verify that at each $((u, j), z, J) \in \mathcal{M}_1^*(\Sigma, M; \beta)$, the system of equations

$$(4.1) \quad D_{J, (j, u)} \bar{\partial}(B, (b, \xi)) = \gamma$$

$$(4.2) \quad D_{J, (j, u)} j_{hol}^k(B, (b, \xi))(z) + \nabla_v(j_{hol}^k(u))(z) = \alpha$$

has a solution $(B, (b, \xi), v) \in T_J \mathcal{J}_\omega \times T_J \mathcal{M}(\Sigma) \times T_u \mathcal{F}_1(\Sigma, M; \beta) \times T_z \Sigma$ for each given data

$$\gamma \in \Omega_{N-1, p}^{(0,1)}(u^* TM), \quad \zeta = (\zeta_0, \zeta_1, \dots, \zeta_k) \in J_{hol}^k(T_z \Sigma, T_{u(z)} M).$$

It will be enough to consider the triple with $b = 0$ and $v = 0$ which we will assume from now on.

We compute the $D_{J, (j, u)} j_{hol}^k(B, (b, \xi))(z)$. It is enough to compute $D_{J, (j, u)} \sigma^l(B, (0, \xi))(z)$ for $l = 0, 1, \dots, k$. We have

$$(4.3) \quad D_{J, (j, u)} \sigma^l(B, (0, \xi))(z) = \pi_{hol} \left((\nabla_{du})^l \xi(z) \right) + \sum_{0 \leq s, t \leq l} B(z) \cdot F_{st}(z) \left((\nabla_{du})^s \xi(z), \nabla^t u(z) \right)$$

where $F_{st}(z)(\cdot, \cdot)$ is some vector-valued monomial, and $B(z)$ is a matrix valued function, both smoothly depending on z . There is no derivative of B in the above formula, because for any l , σ^l is the projection of the tensor $d^l u \in$

$Sym^l(T_z\Sigma, T_{u(x)}M)$ to $Sym_{j,J}^{(l,0)}(T_z\Sigma, T_{u(x)}M)$, and the projection only involves J but not its derivatives. Since u is (j, J) -holomorphic, it also follows that

$$(4.4) \quad \pi_{hol} \left((\nabla_{du})^l \xi(z) \right) = \left(\nabla'_{du} \right)^l \xi(z),$$

where $\nabla'_{du} = \pi_{hol} \nabla_{du} = D_u \partial_{j,J}$. There is a formula for $D_u \partial_{j,J}$ and $D_u \bar{\partial}_{j,J}$ nearby z_0 (see [Si]):

$$(4.5) \quad \begin{aligned} D_u \bar{\partial}_{j,J} \xi &= \bar{\partial} \xi + A(z) \partial \xi + C(z) \xi \\ D_u \partial_{j,J} \xi &= \partial \xi + G(z) \bar{\partial} \xi + H(z) \xi \end{aligned}$$

where $A(z), C(z), G(z), H(z)$ are matrix-valued smooth functions, all vanishing at z_0 .

Now we study the solvability of (4.1) and (4.2) by Fredholm alternative. We regard

$$\Omega_{N-1,p}^{(0,1)}(u^*TM) \times J_{hol}^k(T_z\Sigma, T_{u(z)}M)$$

as a Banach space with the norm

$$\|\cdot\|_{N-1,p} + \sum_{l=1}^k |\cdot|_l$$

where $|\cdot|_l$ is any norm induced by an inner product on the $2n$ -dimensional vector space $Sym_{j,J}^{(l,0)}(T_z\Sigma, T_{u(z)}M) \simeq \mathbb{C}^n$.

We denote the natural pairing

$$\Omega_{N-1,p}^{(0,1)}(u^*TM) \times \left(\Omega_{N-1,p}^{(0,1)}(u^*TM) \right)^* \rightarrow \mathbb{R}$$

by $\langle \cdot, \cdot \rangle$ and the inner product on $Sym_{j,J}^{(l,0)}(T_z\Sigma, T_{u(z)}M)$ by $(\cdot, \cdot)_z$.

Let $(\eta, \zeta) \in \left(\Omega_{N-1,p}^{(0,1)}(u^*TM) \right)^* \times J_{hol}^k(T_z\Sigma, T_{u(z)}M)$ for $\zeta = (\zeta_1, \dots, \zeta_k)$ such that

$$(4.6) \quad \langle D_{J,(j,u)} \bar{\partial}(B, (0, \xi)), \eta \rangle + \sum_{l=1}^k (D_{J,(j,u)} \sigma^l(B, (0, \xi))(z), \zeta_l)_z = 0$$

for all $\xi \in \Omega_{N-1,p}^{(0,1)}(u^*TM)$ and $B \in T_J\mathcal{J}_\omega$. We want to show $(\eta, \zeta) = (0, 0)$. The idea is to change the above equation into

$$\langle D_{J,(j,u)} \bar{\partial}(B, (0, \xi)), \eta \rangle = 0$$

for all ξ and B by judiciously modifying ξ by a Taylor polynomial nearby z , and then use standard techniques in J -holomorphic curve theory to show $\eta = 0$, and after that use Cauchy integral to show $\zeta = 0$. We first deal with $N = k$ case, and later raise the regularity by ellipticity of Cauchy-Riemann equation.

Let $\xi = 0$, then (4.6) becomes

$$\left\langle \frac{1}{2} B \circ du \circ j, \eta \right\rangle = 0.$$

Using the abundance of $B \in T_J\mathcal{J}_\omega$, and that u is a simple J -holomorphic curve, by standard technique (for example [MS]) we get $\eta = 0$ on $\Sigma \setminus \{z_0\}$, namely $\text{supp } \eta \subset \{z_0\}$. Since $\eta \in (W^{k,p})^*$, by the structure theorem of distribution with point support (see [GS]), we have

$$(4.7) \quad \eta = P \left(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}} \right) \delta_{z_0}$$

where δ_{z_0} is the delta function supported at z_0 , and P is a polynomial in two variables with degree $\leq k-1$: this is because the evaluation at a point of the k -th derivative of $W^{k,p}$ maps does not define a continuous functional on $W^{k,p}$.

Let $B = 0$. By (4.3) and (4.7), (4.6) becomes

$$(4.8) \quad \langle D_u \bar{\partial}_{j,J} \xi, \eta \rangle + ((D_{J,(j,u)} j_{hol}^k) \xi(z_0), \zeta)_{z_0} = 0.$$

Since ξ is arbitrary, we can replace ξ by $\xi + \chi(z) h(z, \bar{z})$ in the above identity, where $h(z, \bar{z})$ is a vector-valued polynomial in z and $\bar{z}$, and $\chi(z)$ is a smooth cut-off function equal to 1 in a coordinate neighborhood of z_0 and 0 outside a slightly larger neighborhood, so that $\chi(z) h(z, \bar{z})$ is a well defined and smooth on whole Σ . We want (4.8) becomes $\langle D_u \bar{\partial}_{j,J} \xi, \eta \rangle = 0$ after that replacement. For this purpose the $h(z, \bar{z})$ should satisfy

$$(4.9) \quad \left\langle D_u \bar{\partial}_{j,J} h, P \left(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}} \right) \delta_{z_0} \right\rangle + ((D_{J,(j,u)} j_{hol}^k) h(z_0), \zeta)_{z_0} = - ((D_{J,(j,u)} j_{hol}^k) \xi(z_0, \bar{z}_0), \zeta)_{z_0}$$

After simplification, the above is a differential equation about h :

$$(4.10) \quad Q \left(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}} \right) h(z_0, \bar{z}_0) = w$$

where $Q(s, t)$ is a vector-valued polynomial in two variables s, t , $Q \left(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}} \right)$ acts on $h(z, \bar{z})$ with vector coefficients paired with those of h by inner product, and $w := - ((D_{J,(j,u)} j_{hol}^k) \xi(z_0), \zeta)_{z_0}$ is a constant.

Here comes the crucial observation: when $\zeta_k \neq 0$, the highest degree of s in $Q(s, t)$ is in the term $\zeta_k s^k$. This is because $P(s, t)$ has degree $\leq k-1$ and after integration by parts, $\frac{\partial}{\partial z}$ can fall at $D_u \bar{\partial}_{j,J} h$ of most $(k-1)$ times, and in (4.5)

$$D_u \bar{\partial}_{j,J} h = \bar{\partial} h + A(z) \partial h + C(z) h,$$

where $A(z_0) = 0$. On the other hand, in $(D_{J,(j,u)} j_{hol}^k) h$, by (4.3), (4.4), (4.5), the highest derivative for $\frac{\partial}{\partial z}$ is $\left(\frac{\partial}{\partial z} \right)^k$, and is paired with the coefficient ζ_k in (4.9).

When $\zeta_k \neq 0$, we take $h(z, \bar{z}) = \frac{\zeta_k}{|\zeta_k|^2} \frac{1}{k!} (z - z_0)^k w$, then h solves (4.10). This is because of the following: h is holomorphic nearby z_0 , so we can ignore all terms in $Q \left(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}} \right)$ involving $\frac{\partial}{\partial \bar{z}}$; For the remaining terms in $Q \left(\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}} \right)$, they must be of the form $\left(\frac{\partial}{\partial z} \right)^l$ with $0 \leq l \leq k$, and only $\left(\frac{\partial}{\partial z} \right)^k h(z_0) \neq 0$.

With this h , we reduce the cokernal equation to $\langle D_u \bar{\partial}_{j,J} \xi, \eta \rangle = 0$. Since η is a weak solution of $(D_u \bar{\partial}_{j,J})^* \eta = 0$ on Σ , by ellipticity of the $(D_u \bar{\partial}_{j,J})^*$ operator, the distribution solution η is smooth on Σ (See [Ho]). Since $\eta = 0$ on $\Sigma \setminus \{z_0\}$, $\eta = 0$ on Σ . Then it is not hard to conclude $\zeta = 0$ by Cauchy integral formula as in [OZ] and [Oh]. Therefore the system of equations (4.1) and (4.2) is solvable for any $\eta \in W^{k,p}$ and $\alpha \in J_{hol}^k(T_{z_0} \Sigma, T_{u(z_0)} M)$.

There is one case left: that is when $\zeta_k = 0$. We still need to show $(\eta, \zeta) = (0, 0)$. If $k = 1$, then $\zeta_1 = 0 \Leftrightarrow \zeta = 0$ so it has been done as above. If $k > 1$, we notice that the cokernal equation (4.6) now is the cokernal equation for the section $D\Upsilon_{k-1}$, since the k -th jet is paired with ζ_k there,

$$((D_{J,(j,u)} j_{hol}^k) \xi(z_0), \zeta)_{z_0} = ((D_{J,(j,u)} j_{hol}^{k-1}) \xi(z_0), \zeta)_{z_0}.$$

By induction assumption on k , $D\Upsilon_{k-1}$ has trivial cokernal hence $(\eta, \zeta) = (0, 0)$.

Last we raise the regularity from $W^{k+1,p}$ to $W^{N,p}$, for any $N > k$. For $\eta \in W^{N-1,p} \subset W^{k,p}$, by the above argument we can find a solution $\xi \in W^{k+1,p}$ in (4.1). By elliptic regularity, the solution $\xi \in W^{N,p}$. Therefore (4.1) and (4.2) is solvable in $W^{N,p}$ setting. This finishes induction hence the proof of Theorem.

Remark 1. *In the above proof, the induction starts from $k = 1$. In [OZ], $k = 1$ case was treated in the framework of 1-jet transversality at (u, z_0) where $du(z_0) = 0$. The above proof includes the $k = 1$ case as well, but the way of choosing h does not rely on $du(z_0) = 0$ and applies to any z_0 on Σ .*

Remark 2. *It is crucial that we use the holomorphic k -jet bundle instead of the usual k -jet bundle to get the surjective property of $D\Upsilon_k$. Otherwise, as the usual jet evaluation involves mixed derivatives, given $\zeta_{(k,0)} = 0$ we can not reduce the cokernal equation to the $(k-1)$ case by induction, and when $k = 1$, $\zeta_{(1,0)} = 0$ does not imply $\zeta = 0$. In the case $k = 1$, we can explicitly see why this submersion property fails in the usual 1-jet bundle: for a J -holomorphic curve u with $du(z_0) = 0$, and $\Gamma_1 = (\bar{\partial}, j^k) : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow \mathcal{H}'' \times J^1(\Sigma, M)$, calculations in [OZ] yield*

$$D\Gamma_1(\xi, B) = (D_u \bar{\partial}_{j,J} \xi; D_u \bar{\partial}_{j,J} \xi(z_0), D_u \partial_{j,J} \xi(z_0))$$

therefore there is no solution for $(\eta, \alpha_{(0,1)}, \alpha_{(1,0)})$ if $\eta(z_0) \neq \alpha_{(0,1)}$.

However, if $du(z_0) \neq 0$ then the surjective property still holds in the usual jet bundles. More precisely we have the following

Theorem 2. *At any J -holomorphic curve $((u, j), z_0, J) \in \mathcal{M}_1^*(\Sigma, M; \beta) \subset \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega$ with $du(z_0) \neq 0$, the linearization $D\Gamma_k$ of the section*

$$\Gamma_k = (\bar{\partial}, j^k) : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow \mathcal{H}'' \times J^k(\Sigma, M)$$

is a surjective. Especially the linearization Dj^k of k -jet evaluation

$$j^k : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow J^k(\Sigma, M)$$

at $((u, j), z_0, J)$ is surjective.

Proof. It is enough to show that the cokernal equation

$$\left\langle D_u \bar{\partial}_{j,J} \xi + \frac{1}{2} B \circ du \circ j, \eta \right\rangle + ((D_{J,(j,u)} j^k) \xi(z_0), \zeta)_{z_0} = 0, \quad \text{for all } \xi, B$$

only has trivial solution $(\eta, \zeta) = (0, 0)$. To do this, using standard argument in [MS] we again get $\text{supp } \eta \subset \{z_0\}$. Given $\zeta \in J^k(T_{z_0}\Sigma, T_{u(z_0)}M)$, by Taylor polynomial we can construct a smooth ξ supported in arbitrarily small neighborhood of $z_0 \in \Sigma$, such that $(D_{J,(j,u)} j^k) \xi(z_0) = \zeta$. When $du(z_0) \neq 0$, by linear algebra (namely the abundance of $T_J \mathcal{J}_\omega$) and perturbation method we can construct $B \in T_J \mathcal{J}_\omega$ such that $D_u \bar{\partial}_{j,J} \xi + \frac{1}{2} B \circ du \circ j = 0$ on Σ (see [MS]). So we get from the cokernal equation that $0 + |\zeta|^2 = 0$, i.e. $\zeta = 0$. Let $B = 0$ in the cokernal equation, we get $\langle D_u \bar{\partial}_{j,J} \xi, \eta \rangle = 0$ for all ξ . Then by elliptic regularity we conclude $\eta = 0$ on the whole Σ . $\square$

The following theorem is a direct consequence of Theorem 1 by applying Sard-Smale theorem.

Theorem 3. *Let S be any smooth section of the holomorphic k -jet bundle $J_{hol}^k(\Sigma, M) \rightarrow \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega$. Then the section Υ_k is transversal to the section $(0, S)$. The moduli space*

$$\mathcal{M}^S := (j_{hol}^k)^{-1}(S) \cap \mathcal{M}_1^*(\Sigma, M; \beta) = \Upsilon_k^{-1}(0, S)$$

is a Banach submanifold of codimension $2kn$ in $\mathcal{M}_1^(\Sigma, M; \beta)$. Under the natural projection $\pi : \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega \rightarrow \mathcal{J}_\omega$, there exists $\mathcal{J}_{reg} \subset \mathcal{J}_\omega$ of second category, such that for any $J \in \mathcal{J}_{reg}$, the moduli space $\mathcal{M}_J^S := \mathcal{M}^S \cap \pi^{-1}(J)$ is a smooth manifold in $\mathcal{M}_1^*(\Sigma, M, J; \beta)$, with dimension*

$$\dim \mathcal{M}_J^S = \dim \mathcal{M}_1^*(\Sigma, M, J; \beta) - 2kn,$$

and all the elements in $\mathcal{M}_J^S$ are Fredholm regular.

Remark 3. *In [Oh], the S is the zero section of the holomorphic k -jet bundle $J_{hol}^k(\Sigma, M)$, so $\mathcal{M}_J^S$ is the set of J -holomorphic curves with prescribed ramification degrees at the marked points. The J -holomorphic curves in our moduli space $\mathcal{M}_J^S$ can obey more general constraint S . Similar to [Oh], the theorem also has the version with more than one marked point. Also the constraint S need not to be a full section over the base, but only a closed submanifold in $J_{hol}^k(\Sigma, M)$ whose tangent space projects onto the horizontal distribution of the bundle $J_{hol}^k(\Sigma, M) \rightarrow \mathcal{F}_1(\Sigma, M; \beta) \times \mathcal{J}_\omega$, because the essential part in the proof the theorem is that $D\Upsilon_k|_{(0, S)}$ is surjective.*

The theorem appears to be a good start of studying moduli spaces of J -holomorphic curves satisfying general jet constraints in the holomorphic jet bundle; for example, moduli spaces of J -holomorphic curves with self tangency. Also in [CM], jet constraints from symplectic hypersurfaces were used to get rid of multicovering bubbling spheres. This enables them to define genus zero Gromov-Witten invariants without abstract perturbations.

The above theorem tells that the moduli spaces $\mathcal{M}_J^S$ are well-behaved, and the $\{J_t\}_{0 \leq t \leq 1}$ family version of the above theorem tells that they are cobordant to each other by moduli spaces $\{\mathcal{M}_{J_t}^S\}_{0 \leq t \leq 1}$ for generic path $J_t \subset \mathcal{J}_\omega$. It is interesting to see if the moduli spaces $\mathcal{M}_J^S$ can be used to construct new symplectic invariants.

REFERENCES

- [CM] Cieliebak, K., Mohke, K., “Symplectic hypersurfaces and transversality in Gromov-Witten theory”, Journal of Symplectic Geometry (2008)
- [F] Floer, A., “The unregularized gradient flow of symplectic action”, Comm. Pure Appl. Math. 41 (1988), 775-813
- [GS] Gelfand, I.M., Shilov, G.E., “Generalized Functions”, Vol 2, Academic Press, New York and London, 1968
- [Hir] Hirsch, M., “Differential Topology”, GTM 33, Springer-Verlag, 1976
- [Ho] Hörmander, L., “The Analysis of Linear Differential Operators II”, Compre. Studies in Math. 257, Springer-Verlag, 1983, Berlin
- [HT] M. Hutchings and C. H. Taubes, *Gluing pseudo-holomorphic curves along branched covered cylinders II*, J. Symplectic Geom., 5 (2007), pp. 43–137; math.SG/0705.2074.
- [MS] McDuff D., Salamon, D., “J-Holomorphic Curves and Symplectic Topology”, Colloquim Publications, vol 52, AMS, Providence RI, 2004.
- [Oh] Oh, Y-G, “Higher jet evaluation transversality of J-holomorphic curves”, math.SG/0904.3573
- [OZ] Oh, Y-G., Zhu, K., “Embedding property of J-holomorphic curves in Calabi-Yau manifolds for generic J”, math.SG/0805.3581, Asian J. of Math, Vol 13, No.3, 2009

- [OZ1] Oh, Y.-G., Zhu, K., Floer trajectories with immersed nodes and scale-dependent gluing, submitted, SG/0711.4187
- [Si] Sikorav, J.C., “*Some property of holomorphic curves in almost complex manifolds*”, 165-189, “Holomorphic Curves in Symplectic Geometry”, Audin, M. and Lafontaine, J. ed, Birkhäuser, Basel, 1994
- [Wen] C. Wendl, *Automatic transversality and orbifolds of punctured holomorphic curves in dimension 4*, arXiv:0802.3842v1.

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

E-mail address: `kzhu@math.cuhk.edu.hk`